



PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Wen-Tien CHEN

Serial No.: 09/541,785

Filed: April 3, 2000

For: COMPOSITIONS AND METHODS  
FOR INHIBITION OF CANCER  
INVASION AND ANGIOGENESIS

Examiner: Brumback, B.

Group Art Unit: 1642

Attorney's Docket: 178-295

Dated: July 30, 2002

Assistant Commissioner for Patents  
Washington, D.C. 20231

STATEMENT UNDER 37 C.F.R. §1.804(b)

Sir:

The Rat Hybridomas, E19 and E26, which produce the monoclonal antibody E19 and monoclonal antibody E26, respectively, were deposited with the American Type Culture Collection, Manassas, VA 20110 on May 15, 2001 are the same hybridomas referred to by these names in the present specification as filed on April 3, 2000 and also in the related prior provisional application 60/193,987 filed on April 1, 2000. The above-identified hybridomas and monoclonal antibodies were in Applicants possession prior to filing of each of the above-identified applications.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under §1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Wen-Tien Chen, Ph.D.

Date: July 30, 2002

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Wen-Tien CHEN

Serial No.: 09/541,785

Filed: April 3, 2000

For: COMPOSITIONS AND METHODS  
FOR INHIBITION OF CANCER  
INVASION AND ANGIOGENESIS

Examiner: Brumback, B.

Group Art Unit: 1643

Docket: 178-295

Dated: July 30, 2002

Assistant Commissioner for Patents  
Washington, DC 20231

DECLARATION UNDER 37 CFR 1.132

Sir;

I, Wen-Tien Chen, Ph.D., declare as follows:

1. I received a Ph.D. in Genetics, Cellular and Developmental Biology from Yale University in 1979.
2. Since 1982, I have held faculty positions at a number of accredited universities, including the Howard University Cancer Center at Howard University College of Medicine, the Department of Cell Biology at Georgetown University School of Medicine, and the Lombardi Cancer Center at Georgetown University School of Medicine.

3. Currently, I am a professor in the Department of Medicine/Division of Neoplastic Diseases at the State University of New York at Stony Brook, New York. In addition, I am the director for the Program of Angiogenesis and Metastasis at the Long Island Cancer Center at the State University of New York at Stony Brook.
4. I served as a committee member for the National Institutes of Health in review panels for Biomedical Sciences Study Section from 1984 to 1987 and Cell Biology Study Section from 1989 to 1993.
5. For various periods from 1984 to 2002, I served as an ad hoc committee member for the following: Cancer Center Support Grants, site visits; National Cancer Institute Cancer Preclinical Program Project Review Committee, site visits; National Cancer Institute Special Review Committee, Outstanding Investigator Grants; national Institutes of Health Special Review Committee; RFA De-98-008 Genetic Mechanisms in Oral Cancer; RFA 86-HL-30-L Fibroblast Heterogeneity in Pulmonary Fibrosis; and National Pathology B Study Section, Special Study Section Review.
6. I am an author of over twenty peer-reviewed publications concerning cancer invasion and angiogenesis.

7. I am an expert in the field of cancer invasion and angiogenesis as evidenced by my curriculum vitae, which is attached as Exhibit 1.
8. I am the sole inventor of the above-identified U.S. patent application Serial No: 09/541,785 titled "Compositions and Methods for Inhibition of Cancer Invasion and Angiogenesis."
9. Over one hundred monoclonal antibodies directed against human placental dipeptidyl peptidase IV (DPPIV) have been produced in my laboratory under my direct supervision and control.
10. Only twenty-six of the more than one hundred monoclonal antibodies directed against the DPPIV epitope of the DPPIV-seprase complex were determined by Western Blot analysis to bind to DPPIV with high affinity.
11. I have investigated the ability to inhibit angiogenesis of the twenty-six high affinity monoclonal antibodies directed against the DPPIV-seprase complex described above and three high affinity monoclonal antibodies directed against CD26 available from other sources. These twenty-nine high affinity monoclonal antibodies are listed in Exhibit 2.

12. The twenty-nine monoclonal antibodies shown in Exhibit 2, include the twenty-six high affinity antibodies produced in my laboratory (E1-F102), the 1F7 antibody directed against CD26 and disclosed in Duke-Cohan et al., WO 96/38550 cited by the Examiner, and two commercial preparations also directed against CD26 (202.36 from Ancell and M-A261 from PharMingen).
13. Angiogenesis involves degradation of collagen and cellular invasion. Degradation of gelatin containing denatured collagen is used as one model for angiogenesis. Another model for angiogenesis is invasion of endothelial cells in type 1 collagen and matrix gels.
14. The high affinity monoclonal antibodies listed in Exhibit 2 were tested for their ability to inhibit angiogenesis by examining gelatin degradation and invasion of endothelial cells in type I collagen and matrix gels.
15. Gelatin degradation by the DPPIV-seprase complex was determined by measuring the increase in fluorescence in a mixture of protease complexes and fluorescently labeled gelatin. The experimental protocol for measuring such proteolytic activities of protease complexes is detailed in Exhibit 3. The ability of the high affinity monoclonal antibodies to inhibit

the release of fluorescent-labeled gelatin by immunocaptured DPPIV-seprase complexes was measured. The results are listed in Exhibit 2.


16. Invasion was determined by the inhibition of endothelial cell migration in type I collagen and matrix gels. The experimental protocol for determining invasion of endothelial cells in collagen and matrix gels was described in the legend for Figures 2h-j and Example 4 of the specification of U.S. Serial No. 09/541,785 as originally filed on April 3, 2000.
17. Of the high affinity monoclonal antibodies listed in Exhibit 2, only two monoclonal antibodies (i.e., E19 and E26) exhibited anti-angiogenesis properties as evidenced by inhibition of gelatin degradation and inhibition of endothelial cell migration in type I collagen and matrix gels.
18. Figures 2h-j of U.S. Serial No. 09/541,785 as originally filed on April 3, 2000 demonstrate that the high affinity monoclonal antibody E3 did not inhibit endothelial cell migration. In contrast, Figures 2h-j show that the high affinity monoclonal antibodies E19 or E26 inhibited endothelial cell migration.
19. Thus, only E19 and E26 of the twenty-nine high affinity monoclonal antibodies from over one hundred monoclonal antibodies inhibited

angiogenesis. Accordingly, the ability of the E19 and E26 monoclonal antibodies to inhibit angiogenesis is a rare property among monoclonal antibodies directed against DPPIV.

20. The high affinity monoclonal antibody 1F7 disclosed in Duke-Cohan et al. did not inhibit gelatin degradation and invasion by endothelial cells.

21. The Hartel-Schenk et al. reference (*Eur. J. Biochem.* 1991, 196:369) provides no evidence that the disclosed antibodies directed against DPPIV inhibit angiogenesis. Furthermore, since the ability to inhibit angiogenesis is rare among monoclonal antibodies against DPPIV, there is no reasonable expectation that the monoclonal antibodies disclosed in Hartel-Schenk et al. would inhibit angiogenesis.

I hereby declare that all statements made herein of my own knowledge are true, and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.



Wen-Tien Chen, Ph.D.

Date: July 30, 2002

**THIS PAGE LEFT BLANK**



## **Wen-Tein Chen, Ph.D.**

Professor

Department of Medicine/Division of Neoplastic Diseases

State University of New York

Stony Brook, New York

### **EDUCATION:**

Bachelors of Science degree in 1968 from Tunghai University in Taiwan in field of biology.

Masters of Science degree in 1974 from Northeastern University in Massachusetts in field of embryology.

Doctorate of Philosophy degree in 1979 from Yale University in Connecticut in field of genetics, cellular and developmental biology.

### **POSITIONS AND HONORS:**

- |             |  |
|-------------|--|
| 1974-1979   | Research Assistant, Biology Department, Yale University. (with Dr. J.P. Trinkaus).   |
| 1979-1982   | Postdoctoral Fellow in the area of membrane chemistry, University of California at San Diego. (with Dr. S.J. Singer).                                  |
| 1982-1984   | Director, Cytochemistry Division, Howard University Cancer Center; Assistant Professor, Department of Oncology, Howard University College of Medicine. |
| 1984-1991   | Associate Professor, Department of Cell Biology, Georgetown University School of Medicine.   |
| 1988-1992   | Director, The Cytochemistry & Morphology Core, Lombardi Cancer Center. Georgetown University School of Medicine.                                       |
| 1988-8/1998 | Director, Program of Cellular Control of Invasion and Metastasis, Lombardi Cancer Center, Georgetown University School of Medicine.                    |
| 1992-1993   | Sabbatical in Section of Molecular Biology, NIDR, NIH (Dr. Yoshi Yamada).  |
| 1992-1998   | Professor, Department of Cell Biology, Georgetown University School of Medicine.   |
| 8/1998-     | Professor, Department of Medicine/Division of Neoplastic Diseases, SUNY @ Stony Brook.   |
| 8/1998-     | Director, Program of Angiogenesis and Metastasis, Long Island Cancer Institute. SUNY @ Stony Brook.  |

## **REVIEW PANELS:**

### **(1). Committee Member:**

National Institutes of Health, Cell Biology Study Section (CBY-1), 1989-1993;  
National Institutes of Health, Biomedical Sciences Study Section (BI3), 1984-1987.

### **(2). Ad Hoc Committee Member:**


Cancer Center Support Grants, Site Visits in 2/12-2/14/02; 10/11-10/13/00; 5/31-6/2/2000; 6/29-7/1/98, 10/6-10/8/97;  
NCI Cancer Preclinical Program Project Review Committee, Site Visits in 2/28-3/2/99, 9/16-9/18/96, 1/23-1/25/96;  
NCI Special Review Committee, Outstanding Investigator Grants: 12/9-12/11/92;  
NIH Special Review Committee: 1/31-2/2/99, 3/1991, 3/1990, 3/1987;  
RFA DE-98-008 Genetic Mechanisms in Oral Cancer, 1/31-2/2/1999;  
RFA 86-HL-30-L Fibroblast Heterogeneity in Pulmonary Fibrosis, 7/8-7/9/1987;  
NIH Pathology B Study Section, Special Study Section Review in 4/1984.

## **SELECTED PUBLICATIONS:**

1. Chen, W.-T. 1979. Induction of spreading during fibroblast movement. *J. Cell Biol.* 81:684-691.
2. Chen, W.-T. 1981. Mechanism of retraction of the trailing edge during fibroblast movement. *J. Cell Biol.* 90:187-200.
3. Chen, W.-T. 1981. Surface changes during retraction-induced spreading of fibroblasts. *J. Cell Sci.* 49:1-13.
4. Chen, W.-T., and S.J. Singer. 1980. Fibronectin is not present in the focal adhesions formed between normal cultured fibroblasts and their substrata. *Proc. Natl. Acad. Sci. U.S.A.* 77:7318-7322.
5. Chen, W.-T., and S.J. Singer. 1982. Immunoelectron microscopic studies on the sites of cell-substratum and cell-cell contacts in cultured fibroblasts. *J. Cell Biol.* 95:205-222.
6. Chen, W.-T., K. Olden, B. A. Bernard, F. F. Chu. 1984. Expression of transformation-associated protease(s) that degrade fibronectin at cell contact sites. *J. Cell Biol.* 98:1546-1555.
7. Chen, W.-T., T. Hasegawa, E. Hasegawa, C. Weinstock, and K.M. Yamada. 1985. Development of cell surface linkage complexes in cultured fibroblasts. *J. Cell Biol.* 100: 1103-1114.
8. Chen, W.-T., J.-M. Chen, S.J. Parsons, and J.T. Parsons. 1985. Local degradation of fibronectin at sites of expression of the transforming gene product pp60<sup>src</sup>. *Nature* 316: 156-158.
9. Chen, W.-T., J.-M. Chen, and S.C. Mueller. 1986. Coupled expression and colocalization of 140K cell adhesion molecules, fibronectin, and laminin during morphogenesis and cytodifferentiation of chick lung cells. *J. Cell Biol.* 103: 1073-1090.
10. Chen, J.-M., and W.-T. Chen. 1987. Fibronectin-degrading proteases from the membranes of transformed cells. *Cell* 48: 193-203.
11. Chen, W.-T. 1989. Proteolytic activity of specialized surface protrusions formed at rosette contact sites of transformed cells. *J Exptl. Zool.* 251:167-185.

12. Mueller, S.C., T. Kelly, M. Dai, H. Dai, and W.-T. Chen. 1989. Dynamic cytoskeleton-integrin associations induced by cell binding to immobilized fibronectin. *J. Cell Biol.* 109:3455-3464.
13. Aoyama, A., and W.-T. Chen. 1990. A 170-kDa membrane-bound protease is associated with the expression of invasiveness by human malignant melanoma cells. *Proc. Natl. Acad. Sci. U.S.A.* 87:8296-8300.
14. Mueller, S.C., Y. Yeh, and W.-T. Chen. 1992. Tyrosine phosphorylation of membrane proteins mediates cellular invasion by transformed cells. *J. Cell Biol.* 119:1309-1325.
15. Chen, W.-T. 1992. Membrane proteases: roles in tissue remodeling and tumor invasion. *Current Opinion in Cell Biology.* 4:802-809.
16. Monsky, W.L., T. Kelly, C.-Y. Lin, Y. Yeh, W.G. Stetler-Stevenson, S.C. Mueller, and W.-T. Chen. 1993. Binding and localization of 72 kDa matrix metalloproteinase at cell surface invadopodia. *Cancer Res.* 53: 3159-3164.
17. Kelly, T., S.C. Mueller, Y. Yeh, and W.-T. Chen. 1994. Invadopodia promote proteolysis of a wide variety of extracellular matrix proteins. *J. Cell Physiol.* 158:299-308.
18. Monsky, W.L., C.-Y. Lin, A. Aoyama, T. Kelly, S.K. Akiyama, S.C. Mueller, and W.-T. Chen. 1994. A potential marker protease of invasiveness, seprase, is localized on invadopodia of human malignant melanoma cells. *Cancer Res.* 54: 5702-5710.
19. Chen, W.-T., C.-C. Lee, L. Goldstein, S. Bernier, C. H.L. Liu, C.-Y. Lin, Y. Yeh, W. L. Monsky, T. Kelly, M. Dai, J.-Y. Zhou, and S. C. Mueller. 1994. Membrane proteases as potential diagnostic and therapeutic targets for breast malignancy. *Breast Cancer Research and Treatment.* 31:217-226.
20. Chen, W.-T. 1996. Proteases associated with invadopodia, and their role in degradation of extracellular matrix. *Enzyme Protein* 49: 59-71.
21. Nakahara, H., M. Nomizu, S. K. Akiyama, Y. Yamada, Y. Yeh, and W.-T. Chen. 1996. A mechanism for regulation of melanoma invasion: Ligation of  $\alpha 6 \beta 1$  integrin by laminin G-peptides. *J. Biol. Chem.* 271: 27221-27224.
22. Piñeiro-Sánchez, M., L. A. Goldstein, J. Dodt, L. Howard, Y. Yeh, H. Tran, W. S. Argraves, and W.-T. Chen. 1997. Identification of the 170 kDa melanoma membrane-bound gelatinase (seprase) as a serine integral membrane protease. *J. Biol. Chem.* 272: 7595-7601.
23. Nakahara, H., L. Howard, E. W. Thompson, H. Sato, M. Seiki, Y. Yeh, and W.-T. Chen. 1997. Transmembrane domain mediated membrane type 1-matrix metalloprotease docking to invadopodia is required for cell invasion. *Proc. Natl. Acad. Sci. U. S. A.* 94: 7959-7964.
24. Goldstein, L. A., G. Gherzi, M. Piñeiro-Sánchez, M. Salamone, Y. Yeh, D. Flessate, and W.-T. Chen. 1997. Molecular cloning of seprase: A serine integral membrane protease from human melanoma. *Biochem. Biophys. Acta.* 1361:11-19.
25. Nakahara, H., S.C. Mueller, M. Nomizu, Y. Yamada, and W.-T. Chen. 1998. Activation of  $\beta 1$  integrin signaling stimulates tyrosine phosphorylation of p190RhoGAP and membrane protrusive activities at invadopodia. *J. Biol. Chem. Communication Article.* 273: 9-12.
26. Zukowska-Grojec, Z., E. Karwatowska-Prokopczuk, W. Rose, J. Rone, S. Movafagh, H. Ji, Y. Yeh, W.-T. Chen, H. K. Kleinman, E. Grouzmann, and D. S. Grant. 1999. Neuropeptide Y: A novel angiogenic factor from the sympathetic nerves and endothelium. *Circulation Research* 83(2):187-95.

27. Mueller, S.C., G. Gherzi, S. Akiyama, Q.-Z. Sang, H. Nakahara, L. Howard, M. Piñeiro-Sánchez, Yunyun Yeh, and W.-T. Chen. 1999. A novel protease-docking function of integrin. *J. Biol. Chem.* 274: 24947-24952.
28. Chen, W.-T., and J.-Y. Wang. 1999. Specialized surface protrusions of invasive cells, invadopodia and lamellipodia, have differential MT1-MMP, MMP-2 and TIMP-2 localization. In Conference on Inhibition of Matrix Metalloproteinases: Therapeutic Applications. Annals New York Academy of Science 878: 361-371.
29. Goldstein, L. A., and W.-T. Chen. 2000. Identification of an alternatively spliced seprase mRNA that encodes a novel intracellular isoform. *J. Biol. Chem.* 275: 2554-2559.
30. Zucker, S., J. Cao, and W.-T. Chen. 2000. Critical appraisal of the use of matrix metalloproteinase inhibitors in cancer treatment. *Oncogene*. 19, 6642-6650.
31. Pavlaki, M., Cao, J., Hymowitz, M., Chen, W.-T., Bahou, W., Zucker, S. 2002. A conserved sequence within the propeptide domain of membrane type 1-matrix metalloproteinase (MT1MMP) is critical for function as an intramolecular chaperone. *J. Biol Chem.* 277: 2740-2749
32. Zucker, S., Chen, W.-T. (Editors) 2002. Cell Surface Proteases. Acad. Press, San Diego, CA.
33. Chen, W.-T., T. Kelly, G. Gherzi. 2002. DPPIV, Seprase and Related Serine Peptidases in Multiple Cellular Functions. In Zucker, S., Chen, W.-T. (Editors) Cell Surface Proteases. Acad. Press, San Diego, CA.
34. Gherzi, G., Dong, H., Goldstein, L.A., Yeh, Y., Hakkinen, L., Larjava, H., & Chen, W.-T. 2002. Regulation of fibroblast migration on collagenous matrix by a cell surface peptidase complex. *J. Biol. Chem.* (in press).



**THIS PAGE LEFT BLANK**

**Table 1. Effects of monoclonal antibodies against DPPIV on gelatin-degradation and cellular invasiveness.**

| mAb                    | antibody type | epitope* | Inhibition of gelatin-degradation** | Inhibition of invasion*** |
|------------------------|---------------|----------|-------------------------------------|---------------------------|
| E1                     | rat IgG2a     | DPPIV    | -                                   | -                         |
| E3                     | rat IgG2a     | DPPIV    | -                                   | -                         |
| E5                     | rat IgG1      | DPPIV    | -                                   | -                         |
| E6                     | rat Ig ?      | DPPIV    | -                                   | -                         |
| E7                     | rat IgG2a     | DPPIV    | -                                   | -                         |
| E10                    | rat Ig ?      | DPPIV    | -                                   | -                         |
| E19                    | rat IgG2a     | DPPIV    | +                                   | +                         |
| E20                    | IgM           | DPPIV    | -                                   | -                         |
| E26                    | rat IgG2a     | DPPIV    | +                                   | +                         |
| E30                    | rat IgG2a     | DPPIV    | -                                   | -                         |
| E31                    | rat Ig ?      | DPPIV    | -                                   | -                         |
| E33                    | rat IgG2a     | DPPIV    | -                                   | -                         |
| E96                    | rat Ig ?      | DPPIV    | -                                   | -                         |
| E97                    | rat Ig ?      | DPPIV    | -                                   | -                         |
| E116                   | rat Ig ?      | DPPIV    | -                                   | -                         |
| F2                     | rat IgG2b     | DPPIV    | -                                   | -                         |
| F4                     | rat IgG2a     | DPPIV    | -                                   | -                         |
| F8                     | rat IgG2b     | DPPIV    | -                                   | -                         |
| F9                     | rat IgG2b     | DPPIV    | -                                   | -                         |
| F10                    | rat IgG2b     | DPPIV    | -                                   | -                         |
| F12                    | rat IgG2a     | DPPIV    | -                                   | -                         |
| F20                    | rat IgG2a     | DPPIV    | -                                   | -                         |
| F24                    | rat IgG2b     | DPPIV    | -                                   | -                         |
| F87                    | rat Ig ?      | DPPIV    | -                                   | -                         |
| F95                    | rat Ig ?      | DPPIV    | -                                   | -                         |
| F102                   | rat Ig ?      | DPPIV    | -                                   | -                         |
| 1F7                    | mouse IgG2b   | CD26     | -                                   | -                         |
| 202.36 (Ansell)        | mouse IgG2b   | CD26     | -                                   | -                         |
| M-A261 (BD PharMingen) | mouse IgG1    | CD26     | -                                   | -                         |

\* Epitope was determined by Western immunoblotting.

\*\* Gelatin-degradation was determined by antibody inhibition of the release of fluorescently labeled denatured type I collagen (gelatin) by immunocaptured DPPIV-seprase complexes,

\*\*\* Invasion was determined by the inhibition of endothelial cell invasion into type I collagen gel that overlaid on the monolayer wound model as described in Figure 2h-j of the specification as originally filed.



**THIS PAGE LEFT BLANK**

### **Proteolytic-substrate soluble assay**

The fluorescein labeled gelatin (DQ-gelatin, Molecular Probes) was used as the substrate for the DPPIV-seprase complex in immunoprecipitates. The gelatin was heavily labeled with DQ-fluorescein in which fluorescence was quenched; increase in fluorescent signal indicates cleavage and release of gelatin peptides into solution. Thus, increase in fluorescence was proportional to proteolytic activity. Briefly, 96-well microtiter plates (Costar, Corning, NY) were coated with rabbit anti-rat antibodies (20 µg/ml) and incubated at 37°C for 2 h. Wells were then rinsed with wash buffer (PBS, 0.1% Tween 20), and excess binding sites were blocked with blocking buffer (2% bovine serum albumin in PBS) for 1 h at 37°C. Blocking buffer was removed; wells were coated with anti-human DPPIV mAbs (E3; 10 µg/ml), anti-human seprase mAbs (D8 or D28; 10 µg/ml), or control antibody in PBS and incubated at 37°C for 2 h. Membrane extracts or WGA (wheat germ agglutinant)-enriched protease supernatants derived from HUVEC (human umbilical vein endothelial cells) or MDA-MB-436 human breast carcinoma cells were added and incubated for 1 h at 4°C. The unbound material was removed; wells were washed three times with wash buffer. Gelatin-degrading activity of the immuno-captured DPPIV-seprase complexes were assayed using 100 µl of DQ-gelatin (25 µg/ml in PBS, pH 7.4) with or without small inhibitors or mAbs specific for DPPIV (0.1-10 µg/ml). The change of fluorescent intensity was read at wavelength of 485 (excitation)/538 (emission) using a Microplate Spectrophotometer System (Molecular Devices).